

Binary String Dynamics

Indranil Dasgupta^{1,a} and Ryan Rohm^{1,b}

^a *Boston University, Boston, Ma 02215, USA*

^b *University of North Carolina, Chapel Hill, NC 27599, USA*

In this paper we investigate the dynamical properties of binary cosmic strings [1]. We find extrinsic curvature dependence of the string action and show that kinks on binary strings are eroded while cusps can play a major role in their evolution.

September 26, 1996

¹*e-mail: dgupta@budo.e.bu.edu, rohm@ryan.bu.edu*

1. Introduction

In a recent paper [1] it was shown that the cosmic string solutions of some grand-unified theories have a microscopic structure which is not cylindrically symmetric, but instead is flattened, giving rise to a preferred transverse direction. In this paper we investigate the dynamical properties of these binary cosmic strings. We begin by deriving an effective action for the binary string, using an approximation whereby two worldsheets defined by the cores of the two string components are taken to have constant (spacelike) separation. Describing the string components by the Nambu-Goto action, we then express the result in terms of one worldsheet, with a transverse spacelike vector determining the separation of the component strings. The classical equations obtained in this description display a macroscopic motion close to that of a single string, while the separation vector leads to an extra light degree of freedom for the string, with short-range interactions. The equations are shown to admit a natural geometrical description in terms of the extrinsic geometry of an embedded surface. Because of the difference in the relevant length scales, one may, to first approximation, consider the bulk motion of the binary string to be independent of the local separation vector and consider the dynamics of the separation vector as it evolves on a background geometry defined by a solution to the classical string equations. In this vein it is also possible to consider the effects of kinks, cusps, *etc.*

2. Extrinsic Curvature Dependence of the Action.

The effective action of long cosmic strings can be obtained by integrating out all high energy modes in a background containing a string-like defect [2], [3]. The resulting action has only the collective coordinates corresponding to the space-time positions of the string. The simplest approximations give the well-known Nambu-Goto action [4]

$$S = T \int d\sigma_1 d\sigma_2 \sqrt{-\det D}, \quad (2.1)$$

where T is the string tension (the mass per unit length of the string), $D_{ij} = \frac{\partial x^\mu}{\partial \sigma_i} \frac{\partial x_\mu}{\partial \sigma_j}$; $i, j = 1, 2$. A *binary* string consists of two cosmic strings bound together by a potential (that may be confining) [1]. Its action can be written as the sum of the individual Nambu-Goto actions of the constituent strings plus an interaction term, i.e, $S = S_1 + S_2 - V$. If the interaction V is sufficiently strong to force the ‘spacing’ of the strings to be constant,

the effective action may be reduced to the geometric quantity $S_1 + S_2$ with the additional constraint on the ‘spacing’ of the component strings.

We will first give a more precise meaning to the term ‘separation’. When the two world sheets are *sufficiently flat and parallel*² we can draw at each point p on the world sheet Σ_1 a space-like normal that intersects the world sheet Σ_2 once. This induces a map from Σ_1 to Σ_2 which allows us to parametrize Σ_1 and Σ_2 by the same coordinates σ_1 and σ_2 . Then locally at least,

$$x_{\Sigma_2}^\mu(\sigma_1, \sigma_2) = x_{\Sigma_1}^\mu(\sigma_1, \sigma_2) + a^\mu(\sigma_1, \sigma_2), \quad (2.2)$$

where the ‘separation’ vector a^μ is a section of the normal bundle on Σ_1 at (σ_1, σ_2) . The map exists locally for an interesting region of small perturbations around flat and parallel worldsheets and may admit multiple realizations.

With this parametrization the sum of the areas of the two string world sheets can be effectively reduced to an integral over a single world sheet. However, superior results are obtained by using a world sheet lying ‘midway’ between Σ_1 and Σ_2 as the basis. We shall refer to this as the effective binary string world sheet Σ . Choosing coordinates σ_1 and σ_2 on Σ one can parametrize Σ_1 and Σ_2 by the same coordinates using *two* normal vectors a_1^μ and a_2^μ on Σ . Locally we can define Σ such that the ‘center of mass’ condition $T_1 a_1^\mu + T_2 a_2^\mu = 0$ is valid. Clearly, the ‘spacing’ between the string components can be fixed by constraining the magnitudes of the vectors a_1 and a_2 to be constant. Then the geometric action $S_1 + S_2$ reduces to, $S = \sum T_I M_I$, with $I = 1, 2$ and

$$M_I = \int d\sigma_1 d\sigma_2 \sqrt{-\det \left[\left\{ \frac{\partial(x + a_I)^\mu}{\partial\sigma_i} \right\} \left\{ \frac{\partial(x + a_I)_\mu}{\partial\sigma_j} \right\} \right]}. \quad (2.3)$$

When fluctuations of a_I^μ are small compared to the bulk motion of the binary string, the square roots can be expanded in powers of the derivatives of a_I^μ . To develop the formalism we shall choose a zweibein e_a^μ ($a = 1, 2$) and a basis for the normal space n_k^μ ($k = 3, 4$) on the world sheet Σ . This tetrad satisfies the relations

² A more complete formulation over the bulk of the string motion is irrelevant, and moreover relatively intractable. Local behaviour of the world sheets is well described by small deformations of the flat and parallel world sheets.

$$\begin{aligned}
e_a^\mu e_{b\mu} &= \eta_{ab} , \\
n_k^\mu n_{l\mu} &= -\delta_{kl} , \\
e_a^\mu n_{k\mu} &= 0 .
\end{aligned} \tag{2.4}$$

Note that η_{ab} has Minkowski signature. We will always follow the convention of raising and lowering indices in the tangent bundle by η_{ab} and in the normal bundle by $-\delta_{ij}$. We choose the zweibein by defining, $\partial_a x^\mu = e_a^\mu \sqrt{\partial_b x^\nu \partial_c x_\nu \eta^{bc}}$. The affine connection Γ_{ab}^c and the extrinsic curvature tensor (or the second fundamental form) K_{ab}^k are then defined by, $\partial_a e_b^\mu = \Gamma_{ab}^c e_c^\mu + K_{ab}^k n_k^\mu$.

Expanding a_I^μ as $a_I^\mu = a_I^k n_k^\mu$ one finds that

$$\partial_b n_k^\mu = K_{bk}^a e_a^\mu + A_{bk}^i n_i^\mu , \tag{2.5}$$

where $K_{bk}^a = -K_{bc}^l \delta_{kl} \eta^{ca}$, and A_{bk}^i is a connection on the normal bundle of the world sheet. The curvature F_{abk}^l of the connection A_{bk}^l is given by, $F_{abk}^l = K_{ak}^c K_{bc}^l - K_{bk}^c K_{ac}^l$, and is generally non-zero. A similar statement holds for the curvature of the connection Γ_{ab}^c . The connections A_{bk}^l and Γ_{ab}^c can not therefore be locally gauged away. Using the connection A_{bk}^l we can define a covariant derivative $D_{kb}^i a_I^k$ as

$$D_{kb}^i a_I^k = -D_{ikb} a_I^k = \partial_b a_I^k \delta_k^i - A_{bk}^i a_I^k . \tag{2.6}$$

To expand (2.3), we'll choose the gauge, $\sigma_1 = t, \sigma_2 = \xi$, where t is time and ξ is the length of the string measured from a particular point on the string. Using $T_1 a_1^\mu + T_2 a_2^\mu = 0$, to relate a_1 and a_2 and defining $T = T_1 + T_2$, and $a^k = a_1^k \sqrt{(T_1/T_2)}$ we get

$$\begin{aligned}
S = T \int dt d\xi (1 - v_t^2) & \left[1 - \frac{1}{2} \left\{ \frac{1}{1 - v_t^2} (D_{kt}^i a^k)^2 - (D_{\xi k}^i a^k)^2 + (a^k J_{ak}^a)^2 - 2 \det(a^k J_{bk}^a) \right\} \right] \\
& + \text{higher order} .
\end{aligned} \tag{2.7}$$

Here v_t is the transverse speed of the string and J_{bk}^a is defined by $J_{\xi k}^\xi = K_{\xi k}^\xi, J_{tk}^\xi = K_{tk}^\xi, J_{tk}^t \sqrt{1 - v_t^2} = K_{tk}^t, J_{\xi k}^t \sqrt{1 - v_t^2} = K_{\xi k}^t$. The world sheet indices a and b are contracted using the tensor η^{cd} while the normal indices k and l are contracted using $-\delta^{mn}$. The determinant is taken over the matrix defined by the indices a and b . Note that the only physical velocity is v_t corresponding to the motion of the string perpendicular to itself.

The extrinsic curvature dependence is explicit. The fields a^k live on the world sheet and couple to the extrinsic curvatures. The small parameters of the expansion are the derivatives of a^k and the eigenvalues of the matrices $a^k K_{bk}^a$. The phenomenology resembles that of a thin band with the transverse vector a identified as the ‘width’ of the band. The coupling to the extrinsic curvatures defines the elastic properties of the band associated with bending of the band in its own plane and in a normal plane. Of particular interest are the twists of the band which now constitute a new propagating mode on the string.

3. Twist Modes

The classical equation of motion for the (unadorned) cosmic string with the empirical Nambu-Goto action can be described as follows [5], [6]. The world sheet is described in Minkowski space by $x^\mu(\xi, t)$ with ξ and t being world sheet coordinates. This has the usual reparametrisation invariance $\xi, t \rightarrow \xi', t'$. Choosing the gauge conditions: $t = x^0$, $\left(\frac{\partial x^i}{\partial t}\right)\left(\frac{\partial x^i}{\partial \xi}\right) = 0$, $\left(\frac{\partial x^i}{\partial t}\right)^2 - \left(\frac{\partial x^i}{\partial \xi}\right)^2 = 1$, one obtains the following equation of motion for the spacelike components of $x^\mu(\xi, t)$:

$$\frac{\partial^2 x^i}{\partial t^2} - \frac{\partial^2 x^i}{\partial \xi^2} = 0. \quad (3.1)$$

The solution to (3.1) is

$$x^i(\xi, t) = 1/2[c^i(\xi - t) + d^i(\xi + t)] \quad (3.2)$$

with, $\left(\frac{\partial c^i}{\partial \xi}\right)^2 = \left(\frac{\partial d^i}{\partial \xi}\right)^2 = 1$. Here $\frac{\partial c^i}{\partial \xi}$ and $\frac{\partial d^i}{\partial \xi}$ are components of spacelike three vectors drawn from the origin with tips lying on the Kibble-Turok sphere of unit radius [5].

In the present case, we assume the bulk motion of the binary string to be unaffected by its binary character and look at the equation of motion of a with a given background solution for the bulk motion of the string. Substituting $\frac{\partial x^\mu}{\partial \sigma_a} / |\frac{\partial x^\mu}{\partial \sigma_a}|$ for e_a^μ in the definition of the second fundamental form (with $\sigma_a = \xi, t$) we have, $K_{ab}^l = -(\partial_a e_b^\mu) n_{k\mu} \delta^{kl} = \frac{-\partial_a (\partial_b x^\mu) n_{k\mu} \delta^{kl}}{|\partial_b x^\mu|}$. In the gauge we have chosen, $|\partial_\xi x^\mu| = |\partial_t x^\mu|$. This implies that $K_{\xi t}^l = K_{t\xi}^l = \beta^l$. Using (3.1) we also get $K_{\xi\xi}^l = K_{tt}^l = \alpha^l$. With this form of K_{bk}^a , we obtain the following equation of motion for a^k :

$$\frac{1}{1 - v_t^2} (D_{kt}^i D_{ijt} a^j) - (D_{k\xi}^i D_{ij\xi} a^j) - 2(a^i \alpha_i) \alpha_k Q_1 + 2(a^i \beta_i) \beta_k Q_2 = 0, \quad (3.3)$$

where $Q_1 = \left[\frac{1}{\sqrt{1-v_t^2}} - \frac{1}{2} \left(1 + \frac{1}{\sqrt{1-v_t^2}} \right) \right]$ and $Q_2 = \left[\frac{1}{\sqrt{1-v_t^2}} \right]$. Terms involving α_k and β_k can be dropped when looking for traveling wave solutions of large momentum. One then obtains a free wave equation implying that the coupling to extrinsic curvatures is not relevant for small wavelength excitations. For wavelengths comparable to the scale of the string curvatures, the extrinsic curvatures of the world sheet couple strongly to the a^k . *Thus energy from the bulk motion of the string can be transferred to long wavelength winding or twisting motions of a^k .* Note that the interaction between the two components of the binary string is taken care of by constraining the magnitude of a to be constant. This still leaves an angular coordinate (describing the ‘twist’ of the binary string) free.

The phenomenology of the twist modes can be explored with particular background bulk motions of the string. As an example consider the closed string solution [6]

$$\mathbf{x} = L[\sin(\phi)\cos(\rho)\hat{e}_1 + \cos(\phi)\cos(\rho)\hat{e}_2], \quad (3.4)$$

where, $L = 2\pi \times \text{mass of the string}$, $\phi = \frac{\xi}{L}$, $\rho = \frac{t}{L}$ and $\hat{e}_1$ and $\hat{e}_2$ are orthonormal vectors. The string looks like a collapsing or expanding circle with maximum circumference equaling $\frac{L}{\mu}$, where $\mu = T_1 + T_2$, is the mass per unit length.

For this family of solutions $\beta^l = 0$, $A = 0$. Also $v_t^2 = \sin^2(t/L)$. We define the twist mode θ to be the angle between a and the z axis. The second fundamental form K_{ab}^l has only one non-trivial element, $\alpha^n = -n \cdot \frac{\partial_t^2 x}{|\partial_t x|} = (1/L)\sec(t/L)$ where n is the unit normal to the world sheet and to the z axis. Varying only the angular coordinate θ in action (2.7) we get the equation of motion

$$s^2 \partial_t^2 \theta - \partial_\xi^2 \theta + 2\sin\theta \frac{s^2}{L^2} [s - (1+s)^2], \quad (3.5)$$

where $s = \sec(t/L)$. The explicit time dependence in s causes energy exchange between bulk motion and winding oscillations. The energy exchange is large and is likely to cause some kind of equipartition of energy between curvature oscillations and twist modes. The effect on kinks, as we show now, is even more dramatic.

4. The Fate of Kinks

Kinks are points on the string where the diagonal element ($K_{\xi\xi}^k$) of the second fundamental form is singular. From (3.2) it is clear that kinks arise from discontinuities in $\frac{\partial c^i}{\partial \xi}$ or $\frac{\partial d^i}{\partial \xi}$. Isolated kinks travel to the ‘left’ or ‘right’ at the speed of light. It is therefore

easier to write the kink solution in the light-cone coordinates, $u = \xi + t, v = \xi - t$. The kink solution is proportional to $\delta(u)$ or $\delta(v)$ for a kink travelling in the backward ξ or forward ξ direction. Precisely because of the singularity in the extrinsic curvature at the kinks, the approximation used for deriving the twist mode equations breaks down. However, physically, the thickness of a strongly bound binary string is $\sim \eta^{-1}$, where η^2 is the mass per unit length of the string and the extrinsic curvature at the kinks is less than η . So for the physical kinks, our scheme of approximation remains valid.

When a long wavelength twist mode interacts with a kink we can keep the δ -function approximation for the kink. The transverse speed v_t varies slowly along the string and may be neglected ($1 - v_t^2 \sim 1$). The equation of motion for θ then simplifies to

$$2\frac{\partial^2\theta}{\partial u\partial v} + 6\gamma\eta^2\delta(v\eta)\sin(2\theta) = 0, \quad (4.1)$$

where γ is proportional to the angle of the kink. To first order in θ the equation is linear and the solution is a free wave (with frequency ω) moving opposite to that of the kink and suffering a phase shift after collision with the kink:

$$\theta_0 = A\exp(i\omega u + i6\gamma\frac{\eta}{\omega}\Theta(v\eta)), \quad (4.2)$$

where $\Theta(v\eta)$ is the step function. To the next order the anharmonic term of order A^3 in (4.1) produces scattering. For large amplitudes the small θ approximation fails, but the total power dissipation is easily estimated by noting that η is the only mass scale relevant to the problem. Hence the total power dissipation from a kink is $\sim \eta^2$. The energy in the kink is $\sim \eta$; so the time taken for the kink to disappear is $\sim \eta^{-1}$. This is a very small time scale compared to the expansion times of large string loops ($\sim L$).

We have solved (4.1) numerically for a straight cosmic string with periodic boundary conditions. With a kink angle of 1 radian and a kink size of $4\eta^{-1}$ we simulated the scattering of a sinusoidal twist wave of wavelength $4\eta^{-1}$ and amplitude 2π traveling in a direction opposite to that of the kink. In Figure 1 we plot the ratio of the energy lost by the kink to the energy incident on it as a function of time. The ratio rises to about 25 in the time scale of $4\eta^{-1}$ and oscillates about that value, implying a quick draining of energy from the kink.

In binary strings the presence of the twist modes thus appears to make the kinks very unstable. Numerical simulations indicate that strings have a natural tendency to intercommute and generate kinks [7]. Cosmologically, most of string length at any time is




Figure 1. Energy dissipation from kink vs time due to interaction with a sinusoidal wave of ‘twist’

contained in one long string per horizon. Numerous kinks should exist on the long strings from previous self-intersections. In the case of binary strings the energy of the kinks is continuously and rapidly drained into the twist modes, i.e the binary strings display an ‘elastic’ resistance to developing sharp edges. Consequently, the binary strings are likely to have more cusps on them than ordinary strings. The reason for this is that kinks and cusps are in some sense dual to each other [8]. The vectors $\frac{\partial c^i}{\partial \xi}$ and $\frac{\partial d^i}{\partial \xi}$ in (3.2) describe curves on a unit sphere (the Kibble-Turok sphere). Discontinuities in these curves represent kinks. But one gets cusps if the curves intersect! For each kink smoothed out by winding modes, a corresponding discontinuity in is removed; partial smoothings reduce the jump in the discontinuity. This causes the curves to intersect more, generating more cusps.

The energy loss from cusps and kinks has been examined in the literature [8], [9]. For non-superconducting strings, the loss of energy due to gravitational radiation seems to be comparable in kinky and cuspy strings. For superconducting strings, kinks can radiate large amounts of power. Binary cosmic strings, even if superconducting, are however likely to have no radiating kinks. For light strings, where gravitational effects are small, particle radiation [10] through cusp annihilations can be the dominant method for shrinkage of string loops [11]. In such a case, long binary strings are expected to radiate copiously at all times. Recently however, it has been argued that cusp annihilations may not be so effective in particle production as thought previously [12]. If this is true then binary strings will radiate only by gravitational processes.

Acknowledgements: The work of I.D was supported by the Department of Energy under the grant DE-FG02-91ER40676.

References

- [1] R. Rohm, I. Dasgupta, Phys. Rev. **D 53**, 1827 (1996)
- [2] A. M. Polyakov, Nucl. Phys. **B 268**, 406 (1986).
- [3] P. Orland, Nucl. Phys. **B 428**, 221 (1994).
- [4] D. Forster, Nucl. Phys. **B 81**, 84 (1974); K. Maeda, N. Turok, Phys. Lett. **B 202**, 376 (1988); R. Gregory, Phys. Lett. **B 206**, 199 (1988); P. Orland, Nucl. Phys. **B 428**, 221 (1994).
- [5] T. W. B. Kibble, N. Turok, Phys. Lett. **B 116**, 141 (1982).
- [6] A. Vilenkin, Phys. Reports **121, 5**, 263 (1985).
- [7] E. P. S. Shellard, in *Cosmic Strings - The Current Status*, Proceedings of the Yale Cosmic String Workshop, New Haven, Connecticut, 1988. Edited by F.S.Accetta and L.M.Krauss. (World Scientific, Singapore, 1989) p.25.
- [8] D. Garfinkle, T. Vachaspati, Phys. Rev. **D 36**, 2229 (1987).
- [9] T. Vachaspati, A. Vilenkin, Phys. Rev. **D 31**, 3052 (1985); T. Vachaspati, A. Vilenkin, *ibid.* **35**, 138 (1987).
- [10] R. H. Brandenberger, Nucl. Phys. **B 293**, 812 (1987); P .Bhattacharjee, Phys. Rev. **D 40**, 3968 (1989); R. H. Brandenberger, A. Sornborger, M. Trodden, Phys. Rev. **D 48**, 940 (1993).
- [11] R. H. Brandenberger, A. Davis, M. Hindmarsh, Phys. Lett. **B 263** , 239 (1991).
- [12] R. Gregory, hep-ph/ 9412232.